

# What a retained Tonic Labyrinthine (TLR) Reflex may look like?



- Poor balance
- Weak neck muscles
- Bumps into things
- Motion sickness
- Head on desk, feet wrapped around legs
- Motion sickness
- Visual, speech and auditory difficulties
- Fatigues easily
- Difficulty with sequencing, organisation
- Holds head in hands when writing
- Difficulty with spatial awareness
- Difficult reading
- Poor comprehension
- Slower at reading
- Difficulty with organisation
- Difficulty following directional instructions
- Stiff, jerky movements
- Muscle tone imbalances
- Tight locked muscles
- Poor head control
- Poor core strength
- Fear of heights
- Difficulty judging distance, depth, speed & space
- Forgetful
- Toe walker
- Poor standing posture
- Hunched posture
- Dislikes physical activity
- Difficulties understanding cause and affect
- Difficulty walking up and down stairs
- Slow unco-ordinated movements
- Tends to cross eyes
- Spatial issues
- Poor sense of rhythm
- Difficulties climbing
- Forward tongue
- Speech problems
- Holds heads forward or to the side
- "W" Sitter



[childrenstherapyservices](https://www.instagram.com/childrenstherapyservices)



Childrens Therapy Services

[www.childrenstherapy.online](http://www.childrenstherapy.online)

[info@childrenstherapy.online](mailto:info@childrenstherapy.online)

ASF registered